

Non-Abelian Structures in BSFT and RR couplings*

Emil T.Akhmedov[†]

February 1, 2008

Abstract

In this talk we show that the tachyon annihilation combined with an approximation, in which string theory non-commutativity structure is captured by the algebra of differential operators on space-time, gives a unified point of view on: non-Abelian structures on D -branes; all lowest energy excitations on D -branes; all RR couplings in type II string theory.

1. The understanding of the relation between large N gauge theories and string theories is probably one of the most important problems in the modern theoretical physics [1]. Such a relation implies a very interesting interrelation between the non-Abelian gauge degrees of freedom of open strings and purely geometrical "gravitational" degrees of freedom of closed strings.

Unfortunately so far we have a poor understanding of the relation in question mainly due to the lack of the knowledge of underlying symmetry of string theory. Moreover, it is not clear how the non-Abelian gauge fields emerge in the case of the near coincident D -branes. As well the clear interpretation of the closed strings in open string terms also remains an open question.

Here we would like to address these problems. To explain how we are going to do that let us remind that open string functionals can be considered as operators acting on the space of paths of space-time [2]. This picture appeared to be quite efficient in **CS** String Field Theory (SFT). In fact, it helped in description of tachyon annihilation and D -branes as solitons in SFT [3, 4]. In this talk we will address similar issues using the formalism of Boundary SFT (BSFT) [5, 6]. This formalism turned out to be helpful in verifying the conjectures of [7] as it was demonstrated in [8, 9, 10, 11]. But before dealing with the operators on the space of paths in BSFT we would like to consider approximation of the string functionals by differential operators on space-time¹. We show that the process of the brane annihilation [7] combined with such an approximation leads to the unified description of backgrounds with various D -brane configurations.

The appearance of the differential operators is also natural from SFT point of view. Configuration space of the open string theory is roughly given by the maps of the interval into space-time. If we approximate the strings by the straight lines (classical limit) the configuration space reduces to the space of end-points of the intervals in space-time. Therefore the functionals on configuration space become the functions of two points and could be interpreted as the kernels of the integral operators.

*Extended version of the talk presented at 10th Tohwa International Symposium on String theory which was held in Fukuoka, July 2001.

[†]address:117259, Moscow, B.Chermushkinskaya, 25, ITEP, Russia, email:akhmedov@itep.ru

¹While the consideration of the full non-Abelian structure of BSFT we postpone for the future.

The expansion around diagonal (image of the interval is a point) leads naturally to the differential operators.

Once we define the approximation in which the non-commutative structure of the BSFT can be captured by the algebra of differential operators in space-time, we proceed with the test how the approximation works. Even in such a truncated version of string theory we derive many "stringy" phenomena. We observe that all lowest open string excitations can be obtained as Goldstone type excitations over tachyon profiles corresponding to the creation of D -branes after D -anti- D -brane ($D - \overline{D}$) annihilations. In particular, non-Abelian symmetries on D -brane bound states become obvious. Moreover, we give a unified description of all couplings of the $D - \overline{D}$ -brane lowest excitations to the RR massless fields [12]. We choose to study the RR couplings in this context due to their simple anomalous origin [13]. In fact, they are sensitive to zero modes and, hence, provide a suitable framework for the discussion of the various open string backgrounds in our approximation. These couplings were extensively studied in [14]-[26]. The picture that emerges could be considered as an explicit realization of some proposals from [27].

In an attempt to provide the unified description of the RR gauge field couplings for various background brane configurations it is natural to consider the off-shell interpolation of these backgrounds. The connection with K-theory uncovered in [28, 29] leads to the description of the RR-gauge field couplings in terms of the superconnection [30]. Thus a natural framework for the universal description of RR couplings would be in terms of a superconnection in some universal infinite-dimensional bundle $\mathcal{E}$ over space-time X which provides interpolation between various brane/anti-brane configurations. This infinite-dimensional bundle $\mathcal{E}$ should be naturally connected with the geometry of X and thus it is not very surprising to find the "tautological" bundle with infinite dimensional fiber isomorphic to space of functions on the base manifolds. This bundle has a rich structure connected with the action of the differential operators in the fiber. This is in perfect agreement with the appearance of the differential operators in the explicit description of the elements of the K-homology groups.

This talk is mainly based on the papers [31, 32], but the sections 4 and 7 contain new considerations and formulas as well.

2. In this section we explain how one can capture non-commutative structure of BSFT by the algebra of differential operators on space-time if the classical approximation and UV limit are taken with a suitable regularization.

As a model example consider bosonic sigma-model² on the disk with unspecified boundary conditions:

$$S_{2d} = \int_D d\tilde{X}_\mu d\tilde{X}^\mu, \quad \text{where } \mu = 0, \dots, 25. \quad (1)$$

The action takes extreme values on the configurations:

$$\Delta \tilde{X}_\mu = 0, \quad *d \tilde{X}_\mu|_{\partial D} = 0, \quad (2)$$

where $*d$ is the normal derivative ∂_n on the boundary. Thus on the classical solutions Neumann boundary conditions hold. On the classical solutions the action S_{2d} is equivalent to:

$$S_{2d}^{cl} = \oint_{\partial D} X_\mu * d X^\mu = \oint_{\partial D} \oint_{\partial D} d\theta d\theta' X_\mu(\theta) H(\theta, \theta') X^\mu(\theta'), \quad \text{if } \tilde{X}_\mu|_{\partial D} = X_\mu(\theta) \quad (3)$$

and θ is some parameterization of the boundary. Here

²All considerations of the sections 2,3 and 4 are easy to generalize to the case of superstrings.

$$H = \sum_k |k| e^{ik(\theta - \theta')}$$

is the kernel of the normal derivative operator $*d$ acting on the harmonic functions on the disk.

It is obvious that the theory (3) is equivalent to the following theory with the additional fields P_μ included:

$$I_B = \oint_{\partial D} \left(P_\mu dX^\mu + P_\mu * dP_\mu \right). \quad (4)$$

More precisely, to have the equivalence between the theories (3) and (4) one should not integrate over zero modes of the new fields P_μ in the functional integral (it may be achieved by insertion of the delta function $\delta[P_\mu(0)]$). If we consider X and P as dynamical variables in the quantum mechanics (4) two-point correlation functions of the fields in this theory are given by:

$$\begin{aligned} \langle X^\mu(\theta) X^\nu(\theta') \rangle_{QM} &= \delta_{\mu\nu} G(\theta, \theta') = \delta^{\mu\nu} \sum_{k \neq 0} \frac{1}{|k|} e^{ik(\theta - \theta')}, \\ \langle P_\mu(\theta) P_\nu(\theta') \rangle_{QM} &= \delta^{\mu\nu} \sum_{k \neq 0} \frac{1}{|k|} e^{ik(\theta - \theta')}, \quad \langle X^\mu(\theta) P_\nu(\theta') \rangle_{QM} = \delta_\nu^\mu \sum_{k \neq 0} \frac{1}{k} e^{ik(\theta - \theta')}. \end{aligned} \quad (5)$$

The non-symmetric part of these correlators leads to non-commutativity of the operators P and X :

$$[X^\mu(\theta), X^\nu(\theta')] = 0, \quad [P_\mu(\theta), P_\nu(\theta')] = 0, \quad [P_\mu(\theta), X^\nu(\theta')] \propto \delta_\mu^\nu \delta(\theta - \theta'). \quad (6)$$

The symmetric part of the correlation functions is given by the standard Green function:

$$G(\theta, \theta') = \log |e^{i\theta} - e^{i\theta'}|^2 \quad (7)$$

and is singular when the points coincide. This leads to the necessity to regularize the theory. Let us suppose that our regularization prescription modifies the metric on the boundary as follows:

$$\begin{aligned} ||\theta - \theta'|| &= |\theta - \theta'| \quad \text{when} \quad |\theta - \theta'| \gg l_R \\ ||\theta - \theta'|| &= 1 \quad \text{when} \quad |\theta - \theta'| \ll l_R, \end{aligned} \quad (8)$$

where l_R is a characteristic regularization length. With this prescription we get for modified Green function $G_R(0) = 0$. Note that this regularization does not modify the "non-commutative" part of the correlators. Thus we would like to conclude that if we are interested in the small distance correlation functions only "non-commutative" parts survive. In another words, with the regularization (8) correlation functions at small distances are dominated by the first term in the lagrangian (4). One would like to propose that this is the way matrix degrees of freedom show up in the sigma model approach. To show this fact it would be interesting to extract relevant degrees of freedom explicitly. This is done below.

3. In this section we show that in our approximation string vertex operators can be represented as differential operators on space-time or more generally on some auxiliary space. To see this let us start with Dirichlet boundary condition $\tilde{X}|_{\partial D} = X(\theta) = Y^1$. It is easy to construct the boundary

operator in conformal field theory that shifts the Dirichlet boundary conditions $\tilde{X}|_{\partial D} = Y^1$ to a new boundary condition $\tilde{X}|_{\partial D} = Y^1 + \Delta X(\theta)$:

$$V_{\Delta X} = e^{-\int d\theta \partial_n X(\theta) \Delta X(\theta)}. \quad (9)$$

In order to realize the process when in the beginning the string ends on the D -brane $\tilde{X}|_{\partial D} = Y^1$ then jumps to D -brane $\tilde{X}|_{\partial D} = Y^2$ and then at the end jumps back to the D -brane $\tilde{X}|_{\partial D} = Y^1$ we take the "shift"-function in the form of step-function:

$$\Delta X = (Y^1 - Y^2) \epsilon(\theta|\theta_1, \theta_2), \quad \text{where} \quad (10)$$

$$\begin{aligned} \epsilon(\theta|\theta_1, \theta_2) &= 1 & \theta_1 < \theta < \theta_2 \\ \epsilon(\theta|\theta_1, \theta_2) &= 0 & \text{otherwise.} \end{aligned}$$

Then using the decomposition of the scalar field $X = X_+ + X_-$ (on the boundary of the disc) in terms of positive and negative frequency modes

$$X^\mu(\theta) = X_+^\mu(\theta) + X_-^\mu(\theta) = \left(\frac{1}{2} X_0^\mu + \sum_{k>0} X_k^\mu e^{ik\theta} \right) + \left(\frac{1}{2} X_0^\mu + \sum_{k<0} X_k^\mu e^{ik\theta} \right) \quad (11)$$

we could represent $V_{\Delta X}$ operator as the product of two "jump"-operators:

$$\begin{aligned} V_{\Delta X} &= \exp \left\{ \int_{\theta_2}^{\theta_1} d\theta (Y^1 - Y^2) \partial_n X \right\} = \exp \left\{ \int_{\theta_2}^{\theta_1} d\theta (Y^1 - Y^2) [\partial_t X_+ - \partial_t X_-] \right\} = \\ &= \exp \left\{ [X_+(\theta_1) - X_-(\theta_1)] (Y^1 - Y^2) \right\} \exp \left\{ -[X_+(\theta_2) - X_-(\theta_2)] (Y^1 - Y^2) \right\}, \end{aligned} \quad (12)$$

where $\partial_t = \partial_\theta$ is the tangential derivative to the boundary and we used the classical equations of motion here. Therefore, one has the following expression for the "jump"-operator from Y^i to Y^j :

$$V^{ij} [X(\theta)] = e^{\frac{(Y^i - Y^j)}{\alpha'} P(\theta)} = e^{\frac{(Y^i - Y^j)}{\alpha'} \frac{\delta}{\delta X(\theta)}}, \quad \text{where} \quad P(\theta) = X_+(\theta) - X_-(\theta). \quad (13)$$

This operator corresponds to the excitation with the mass defined by its conformal dimension:

$$m^2 = |Y^i - Y^j|^2 / (\alpha')^2, \quad (14)$$

which is in accordance with the expectations ($m^2 \propto \text{length}^2$) for the mass of the non-Abelian part of the tachyon. In order to get the operators describing the general states from the spectrum one should multiply (13) by the polynomial of the derivatives of X and the standard exponential factors e^{ipx} responsible for the non-zero momentum along the D -brane.

What is most important for us is that such operators as (13) respect the D -brane state given under the string functional integral by the composition of $\delta(X - Y^i)$ where $i = 1, \dots, k$ in the case if there are k D -branes (see below). As well V^{ij} generate the $gl(k)$ algebra when they act on the system of δ -functions in question [32].

4. In this section we show how non-Abelian degrees of freedom on D -branes appear after $D - \overline{D}$ -brane annihilation if we approximate string functionals by differential operators on an auxiliary space.

Consider the annihilation of infinite number of $D25$ -branes into k Dp -branes. In our approximation the lowest excitations of the $D25$ -branes are described by the following generating functional:

$$Z_{D25} = \left\langle \int \mathbb{D}y^\mu \mathbb{D}p_\mu \mathbb{D}c^\mu \mathbb{D}\bar{c}_\mu \exp \left\{ - \oint_{\partial D} d\theta \left[i p_\mu \partial_t y^\mu + i \bar{c}_\mu c^\mu + \right. \right. \right. \\ \left. \left. \left. + i A_\mu \left(X | y, p \right) \partial_t X^\mu + T \left(X | y, p \right) \right] \right\} \right\rangle_{BS}, \quad (15)$$

where the averaging $\langle \dots \rangle_{BS}$ is taken with the bosonic string functional integral with the action (1). In this formula we stressed that string fields are differential operators on an auxiliary space Y , parameterized by the coordinates y^μ . In fact, when there is infinite number of $D25$ -branes the space on which Chan-Paton indexes act becomes more like a Hilbert space of functions on a manifold Y rather than some finite dimensional vector space. Hence, string functionals are operators acting on this Hilbert space, i.e. they are functions on T^*Y . In "tautological" situation Y coincides with space-time X . It is this situation we considered above and going to consider below.

The integral over the fermions $\bar{c}$ and c gives the determinant of the symplectic form on T^*Y [37]. When $X = Y$ such fermions have a natural origin in superstring theory as boundary values of NSR fermions (superpartners of X) and their conjugates (superpartners of P). While in bosonic string we have to insert them by hand according to [37].

To describe the process of the annihilation of $D25$ -brane into k Dp -branes one has to consider the off-shell tachyon profile as follows [31, 32]:

$$T \left(X | y, p \right) \propto \frac{1}{t} |X^\mu - y^\mu|^2 + \frac{1}{s_1} |W(y^a)|^2 + \frac{1}{s_2} |p_\mu|^2, \quad (16)$$

where $W(y^a)$ is a polynomial of y^a , $a = p + 1, \dots, 25$ of degree k on zeros of which we are going to localize the Dp -branes; t , s_1 and s_2 are some coupling constants. As is argued in [9] the coupling constant t flows to 0 in the IR. Hence, the first term in (16) describes the localization of space Y on space-time X . The second term in (16) describes localization on the Dp -branes, while the reason of the third one is explained in the sections 5 and 6, below.

Let us take the parameters $t, s_1, s_2 \rightarrow 0$. In case if $A_\mu = 0$ and all zeros of W are separated we obtain [32]:

$$\lim_{t, s \rightarrow 0} \int \mathbb{D}y \mathbb{D}p \mathbb{D}c \mathbb{D}\bar{c} \exp \left\{ - \oint_{\partial D} d\theta \left[T \left(X | y, p \right) + i p \partial_t y + i \bar{c} c \right] \right\} \propto \sum_{i=1}^k \delta(X_a - Y_a^i) \quad (17)$$

under the string functional integral. Here Y^i are zeros of W . Thus, in this case we obtain just k Dp -branes without any excitations on them.

Now we can deform the background in such a way that the Dp -brane system (17) is respected in the sense that Dp -brane positions are respected. First, we can consider in (15) $A_\mu(X|y, p) = \left[A_m^g(X^m) \ T^g(y^a, p_a), 0, \dots, 0 \right]$, where $m = 0, \dots, p$ enumerate directions along the Dp -branes, $a = p + 1, \dots, 25$ enumerate directions transversal to the branes ($\mu = (m, a)$) and

$$\left[T^{g_1}, \ T^{g_2} \right] = f^{g_1 g_2 g_3} T^{g_3}, \quad g = 1, \dots, k^2, \quad (18)$$

i.e. T^g generate the $U(k)$ algebra. The relation of such generators to the vertex operators V^{ij} , $i = 1, \dots, k$ (13) is described in the section 3 of [32]. As well for such a choice of A_μ the Dp -brane system (17) is respected, i.e. mapped under the action of the corresponding vertex operators onto itself. Note that after the limit $s_1, s_2 \rightarrow \infty$ the group of symplectomorphisms acting on T^*Y is broken to $U(k)$ [32].

Second, we can shift X_a via the following operator (we use here (9)-(13)):

$$\oint_{\partial D} d\theta \tilde{\Phi}_a^{\tilde{g}}(X^m) T^{\tilde{g}}(y, p) \frac{\delta}{\delta X^a} = \oint_{\partial D} d\theta \tilde{\Phi}_a^{\tilde{g}}(X^m) T^{\tilde{g}}(y, p) \partial_n X^a, \quad (19)$$

for some real functions $\tilde{\Phi}^g$. In this formula we included only generators $T^{\tilde{g}}$, $\tilde{g} = 1, \dots, k^2 - k$ of $U(k)$, which do not belong to its Cartan subalgebra. This is done to respect Dp -brane system (17).

Now after the integration over y, p, c and $\bar{c}$ via localization procedure [37], we obtain:

$$\lim_{s, t \rightarrow 0} Z_{D25} = \left\langle \text{Tr P exp} \left\{ \oint_{\partial D} d\theta \left[i \hat{A}_m(X^l) \partial_t X^m + \hat{\Phi}_a(X^l) \partial_n X^a \right] \right\} \right\rangle_{BS}, \quad (20)$$

where

$$\hat{A}(X^m) = A^g(X^m) t^g, \quad \hat{\Phi}(X^m) = \tilde{\Phi}^{\tilde{g}}(X^m) t^{\tilde{g}} + \text{diag}(Y^1, \dots, Y^k) \quad (21)$$

and $t^g, t^{\tilde{g}}$ are the matrix generators of $U(k)$ in the fundamental representation corresponding to T^g and $T^{\tilde{g}}$ respectively. As well the trace "Tr" in (20) is taken in the fundamental representation of $U(k)$. Hence, if all Y^i coincide the full non-Abelian $U(k)$ symmetry is restored, while if some of Y^i are separated the symmetry is broken down to a subgroup of $U(k)$. In conclusion, this formula already describes massless excitations of k Dp -branes rather than of infinite number of $D25$ -branes.

5. Therefore, in our approximation we can give a geometric unified point of view on Chan-Paton (non-Abelian) structures in BSFT. Let us describe as well the unified approach to all open string massless degrees of freedom within our approximation. We are going to do this on the example of RR couplings to D -branes in type II superstring theory. In this section we just set the notations.

In particular, we describe the RR gauge field couplings with open strings for the backgrounds with infinite number of $D9 - \overline{D9}$ -branes and the low dimensional brane backgrounds arising in the process of the annihilation. The general coupling of the type IIB $D9 - \overline{D9}$ -brane fields with the RR fields is given by

$$S_{RR} = Q_9 \int_{X^{(10)}} \left[C_{RR} \wedge \text{Ch}(\mathbb{A}) \right]_{top}. \quad (22)$$

Here Q_9 is the elementary $D9$ - and $\overline{D9}$ -brane charge, $X^{(10)}$ is ten-dimensional flat target space, C_{RR} is defined as the sum of the RR gauge fields of the given parity (even in IIB theory):

$$C_{RR} = \sum_k C_{(2k)}, \quad C_{(2k)} = C_{\mu_1 \dots \mu_{2k}}(x) dx^{\mu_1} \wedge \dots \wedge dx^{\mu_{2k}} \quad (23)$$

and the subscript *top* means that we should integrate the top-dimensional differential form over $X^{(10)}$; x^μ , $\mu = 0, \dots, 9$ are coordinates on $X^{(10)}$. We use the normalization of RR gauge fields consistent with the standard definition of the superconnection used below.

$\text{Ch}(\mathbb{A})$ is the Chern character of the superconnection $\mathbb{A}$ [30], which is constructed from the open string modes. In fact, the gauge bundles on $D9$ - and $\overline{D9}$ -branes may be combined in a $\mathbb{Z}_2$ -graded bundle with the superconnection defined in terms of the gauge field A_μ on the $D9$ -brane, the gauge field $\tilde{A}_\mu$ on the $\overline{D9}$ -brane and the complex tachyon field T corresponding to the lowest energy excitation of the strings stretched between $D9$ - and $\overline{D9}$ -branes:

$$\mathbb{A} = \begin{pmatrix} dx^\mu \nabla_\mu^+ & T \\ \bar{T} & dx^\mu \nabla_\mu^- \end{pmatrix},$$

$$\nabla_\mu^+ = \partial_\mu + i A_\mu, \quad \text{and} \quad \nabla_\mu^- = \partial_\mu + i \tilde{A}_\mu. \quad (24)$$

The Chern character of the superconnection is defined by the standard formula:

$$\text{Ch}(\mathbb{A}) = \text{Str} e^{-\frac{\mathbb{A}^2}{2\pi}}. \quad (25)$$

Here we use the supertrace "Str" on the matrices acting on $\mathbb{Z}_2$ -graded vector spaces, which is defined as follows. Let $V = V_+ \oplus V_-$ be a $\mathbb{Z}_2$ -graded vector space and:

$$\mathcal{O} : V \rightarrow V, \quad \mathcal{O} = \begin{pmatrix} \mathcal{O}_{++} & \mathcal{O}_{+-} \\ \mathcal{O}_{-+} & \mathcal{O}_{--} \end{pmatrix}. \quad (26)$$

Here V_+ is a bundle on $D9$ -brane and V_- is that on $\overline{D9}$ -brane. Then the supertrace of $\mathcal{O}$ is defined as:

$$\text{Str}_V \mathcal{O} = \text{Tr}_V \tau \mathcal{O} = \text{Tr}_{V_+} \mathcal{O}_{++} - \text{Tr}_{V_-} \mathcal{O}_{--}, \quad (27)$$

where τ is an operator defining $\mathbb{Z}_2$ -structure. In our case the vector space V is infinite-dimensional space of functions on an auxiliary space Y . Hence, the traces "Tr" contain integrals over the phase space T^*Y : $\text{Tr}_{\mathcal{H}} \dots \propto \int dp dy$, where $\mathcal{H} \in V$ is the Hilbert space of functions on Y .

Note that to have a well defined trace $\text{Tr}_{\mathcal{H}}$ the operators should have special property (to be of the trace class). A particular class of such operators is given by the expressions $F(p, y) \propto \delta^{(10)}(p) f(y)$. It's trace is naturally represented by the integral over Y rather than over T^*Y . This type of operators has a simple qualitative interpretation in terms of string backgrounds. Let us symbolically denote open string by the matrix $K_{y_1, y_2}(x)$ (or more exactly integral operators $\hat{K}$ with the kernels $K(x|y_1, y_2)$ acting on the space of functions on Y) where possible end points of the strings are enumerated by the indexes " y_1 " and " y_2 ". The trace invariants naturally correspond to the open strings with identified ends and thus may be considered as closed strings. Now let us consider the trace of the operator $\hat{K}$ with additional insertion of the operator $\delta(p)$ which projects on the states with zero eigenvalue of p . In case if $X = Y$ it is easy to see that:

$$\delta(p) = |p=0\rangle \langle p=0| = \left(\sum_x |x\rangle \right) \left(\sum_{x'} \langle x'| \right) \quad (28)$$

(note that the identity operator is $1 = \sum_x |x\rangle \langle x|$) and we have:

$$\text{Tr}_{\mathcal{H}} \delta(p) \hat{K} \propto \sum_{x, x'} K(x, x') \quad (29)$$

We see that in the presence of the operator $\delta(p)$ we should sum over the positions of the "ends" of the string independently. Thus its insertion may be interpreted as the creation of the D -brane filling whole space-time.

We have replaced the consideration of the infinite number of D -branes by the consideration of the infinite-dimensional vector bundles over the space-time X with the fiber — the space of functions on the copy Y of X . The Lie algebra of the differential operators acts naturally in the fibers of the bundle. Note that one may consider infinite dimensional bundles with the fiber given by the sections of a finite dimensional (super)-bundle. Below we encounter examples with the space of sections of spin bundles as a fiber. Having in mind such more general cases treated below we consider Clifford algebra for the total space $X \otimes T^*Y$:

$$\begin{aligned} \{\gamma_\mu, \gamma_\nu\} &= \delta_{\mu\nu} & \{\Gamma_\mu, \Gamma_\nu\} &= \delta_{\mu\nu} & \{\hat{\Gamma}^\mu, \hat{\Gamma}^\nu\} &= \delta^{\mu\nu} \\ \{\Gamma_\mu, \gamma_\nu\} &= \{\hat{\Gamma}^\mu, \Gamma_\nu\} & \{\hat{\Gamma}^\mu, \gamma_\nu\} &= 0. \end{aligned} \quad (30)$$

The gamma matrices are defined with respect to the explicit coordinates as follows: $(x^\mu, y^\nu, p_\rho) \leftrightarrow (\gamma_\mu, \Gamma_\nu, \hat{\Gamma}^\rho)$. Note that Γ and $\hat{\Gamma}$ (as well as γ) have a natural interpretation as zero modes of superpartners of y and p (X) in superstring analog of (15).

6. Now we give an explicit construction of the superconnection in the infinite-dimensional bundle in question which leads to the desired description of RR-couplings. We would like to find such an $\mathbb{A}$ which after the annihilation of the infinite number of $D9 - \overline{D9}$ -branes leads to one $D9$ -brane. Similarly to (16) we have to obtain something like³:

$$\left| T \begin{pmatrix} x & y, p \end{pmatrix} \right|^2 \propto \frac{1}{t} |x^\mu - y^\mu|^2 + \frac{1}{s} |p_\mu|^2. \quad (31)$$

In fact, the first term is responsible for localization of Y on X while the second one leads eventually the trace class operator from $\text{Ch}(\mathbb{A})$. Below we consider localizers like W in (16) when we study creation of Dp -branes with $p < 9$.

At the same time according to [29] we have to consider such a T which gives a non-trivial map from one chirality spinor bundles on $X \otimes T^*Y$ to another. Hence, we have to embed part of the "Chan-Paton" indexes of T into spinor bundles on $X \otimes T^*Y$. This means that we take $V = V_+ \oplus V_-$ such that $V_\pm = s_\pm \otimes S_\pm \otimes \hat{S}_\pm \otimes \mathcal{H}$ and $s_\pm, S_\pm$ and $\hat{S}_\pm$ are spinor bundles on X, Y and on the fibers of T^*Y , respectively. Thus we take the value for the tachyon as follows:

$$T \begin{pmatrix} x & y, p \end{pmatrix} = \frac{1}{\sqrt{t}} \sigma_\mu (x^\mu - y^\mu) + \frac{1}{\sqrt{s}} \hat{\Sigma}^\mu p_\mu. \quad (32)$$

In this formula σ and $\hat{\Sigma}$ are σ -matrixes corresponding to γ -matrixes from (30).

As well we choose trivial connections on both $D9$ - and $\overline{D9}$ -branes $\nabla = \nabla^+ + \nabla^- = d$. Hence, the corresponding superconnection is

$$\mathbb{A}_{s,t} = d + \frac{1}{\sqrt{t}} \gamma_\mu (x^\mu - y^\mu) + \frac{1}{\sqrt{s}} \hat{\Gamma}^\mu p_\mu \quad (33)$$

This superconnection takes values in the space of differential operators acting in the auxiliary space of sections of spinor bundle on $X \otimes T^*Y$. The $\mathbb{Z}_2$ -structure in this case is defined by the chirality.

³Note that the tachyon vertex on the $D - \overline{D}$ -brane systems in type II superstring is $\oint |T|^2$ rather than $\oint T$ like in bosonic string [10].

We mean that with our choice of V the Z_2 -grading operator is defined by $\tau = \gamma^{11} \otimes \Gamma^{11} \otimes \widehat{\Gamma}^{11}$, where γ^{11} , etc. are ten-dimensional analogs of γ^5 -matrix in four dimensions.

The square of (33) is given by the expression:

$$\mathbb{A}_{s,t}^2 = \frac{1}{t} |x^\mu - y^\mu|^2 + \frac{1}{s} |p_\mu|^2 + \frac{1}{\sqrt{st}} \gamma_\mu \widehat{\Gamma}^\mu + \frac{1}{\sqrt{t}} dx^\mu \gamma_\mu. \quad (34)$$

The resulting Chern character we are looking for is defined as:

$$\text{Ch}(\mathbb{A}) = \lim_{t \rightarrow 0} \lim_{s \rightarrow 0} \text{Ch}(\mathbb{A}_{s,t}) \quad (35)$$

Note that here "Str" is taken over the representation of the full Clifford algebra (30) and includes $\text{Tr}_{\mathcal{H}}$ as well, i.e. with our choice of V we have:

$$\text{Str} \dots = \text{sp } \gamma^{11} \text{ Sp } \Gamma^{11} \widehat{\text{Sp}} \widehat{\Gamma}^{11} \text{Tr}_{\mathcal{H}} \dots,$$

where by "sp", etc. we define the traces in the spinor part of the bundle V to distinguish them from "Tr" over $\mathcal{H}$.

Thus, using the identities:

$$\text{sp} \left(\gamma^{11} \gamma_{\mu_1} \dots \gamma_{\mu_{10}} \right) = (2i)^5 \epsilon_{\mu_1 \dots \mu_{10}} \quad (36)$$

and

$$\lim_{t \rightarrow 0} \frac{1}{(\pi t)^5} \exp \left\{ -\frac{|x - y|^2}{t} \right\} = \delta^{(10)}(x - y), \quad (37)$$

(and similarly for p) we obtain:

$$\text{Ch}(\mathbb{A}) = \text{Tr}_{\mathcal{H}} \delta^{(10)}(p_\mu) \delta^{(10)}(x^\mu - y^\mu) = 1. \quad (38)$$

Which means that we have reproduced the correct coupling with the top-degree RR-form for the $D9$ -brane filling whole space-time:

$$S_{RR} = Q_9 \int_{X^{(10)}} C_{(10)}(x).$$

This can be considered as the proof that we have made a proper choice of the background $\mathbb{A}$ (33).

7. Now let us give a general construction of the RR gauge field couplings for less trivial D -brane backgrounds. We start with a fixed infinite-dimensional bundle over space-time with the fiber identified with the space of sections of the spinor bundle over the base space times the two-dimensional vector space. We start with the construction of a superconnection corresponding to parallel k_+ $D7$ -branes and k_- $\overline{D7}$ -branes. Let y^a , $a = 8, 9$ be coordinates transversal to the $D7$ -branes; $W(y^a)$ is a function defined by the condition that its critical points are the positions of $D7$ - and $\overline{D7}$ -branes in the plane (y_9, y_8) : the sign of the Hessian $\partial^2 W$ defines the sign of the RR charge. Consider now the superconnection:

$$\mathbb{A}_{s,t} = d + \frac{1}{\sqrt{t}} \gamma_\mu (x^\mu - y^\mu) + \frac{1}{\sqrt{s_1}} \Gamma_a \partial^a W(y) + \frac{1}{\sqrt{s_2}} \widehat{\Gamma}^\mu p_\mu \quad (39)$$

In this formula the additional two-dimensional vector space is interpreted as the representation of the Clifford algebra of the cotangent space to the transversal space to the $D7$ -branes (the notations are in agreement with (30)). Note that this representation is very close to the expression that enter the Supersymmetric Quantum Mechanic interpretation of Morse theory [36].⁴ In fact, the calculation of the Chern character for this connection in the limit $t, s_1, s_2 \rightarrow 0$ gives the expression with the insertion of the projector in the y -integral:

$$\det \left[\partial^a \partial_b W \right] \delta^{(2)} \left[\partial_a W(y) \right] \quad (40)$$

along with $\delta^{(10)}(x - y)$ and $\delta^{(10)}(p)$. Trivial calculation of $\text{Tr}_{\mathcal{H}}$ reduces the full expression to the Chern character form representing k_+ $D7$ -branes and k_- $\overline{D7}$ -branes ($k_{\pm}$ are the numbers of critical points of $W(y)$ with the positive and negative indexes):

$$S_{RR} = Q_7 \int_{X^{(10)}} C_{(8)}(x^\mu) \left\{ \sum_{i=1}^{k_+} \delta(x^a - Y_i^{a+}) - \sum_{j=1}^{k_-} \delta(x^a - Y_j^{a-}) \right\},$$

where Q_7 is the elementary $D7$ -brane charge and $Y^\pm$ are critical points of $W(y)$ with the positive and negative indexes.

Let us consider now deformations of (39) which correspond to the massless excitations on the $D7$ -branes in case if there are no $\overline{D7}$ -branes after the annihilation. We are interested in deformations which preserve the $D7$ -brane system. There are many gauge equivalent ways to deform the superconnection (39), which are generated by the factor of symplectomorphisms acting on T^*Y over $U(k)$. As well there are many deformations which correspond to massive excitations over the chosen tachyon background and lead to arbitrary (not necessary parallel) configuration of $D7$ -branes. In this talk we are not going to study the latter kind of deformations.

The simplest deformations which correspond to the massless modes are described by:

$$\begin{aligned} \mathbb{A}_{s,t} = d + \frac{1}{\sqrt{t}} \gamma_m (x^m - y^m) + \frac{1}{\sqrt{t}} \gamma_a \left\{ x^a - y^a - \tilde{\Phi}^a(y^l | y^a, p_a) \right\} + \frac{1}{\sqrt{s_1}} \Gamma_a \partial^a W(y) + \\ + \frac{1}{\sqrt{s_2}} \hat{\Gamma}^m \left\{ p_m + i A_m(y^l | y^a, p_a) \right\} + \frac{1}{\sqrt{s_2}} \hat{\Gamma}^a p_a, \end{aligned} \quad (41)$$

where $A_m(y^l | y^a, p_a)$, $\tilde{\Phi}_a(y^l | y^a, p_a)$ are defined as in the section 4 with $m, l = 0, \dots, 7$ and $a = 8, 9$; y^m are coordinates along the directions parallel to the $D7$ -brane world-volume.

The square of this superconnection is

$$\begin{aligned} \mathbb{A}_{s,t}^2 = \frac{1}{2s_2} \left[\nabla_m, \nabla_n \right] \hat{\Gamma}^m \hat{\Gamma}^n + \frac{1}{\sqrt{t}} dx^\mu \gamma_\mu + \frac{1}{\sqrt{s_2 t}} \left[\nabla_m, \tilde{\Phi}^a \right] \gamma_a \hat{\Gamma}^m + \\ + \frac{1}{2t} \left[\tilde{\Phi}^a, \tilde{\Phi}^b \right] \gamma_a \gamma_b + \frac{1}{t} \left| x^a - y^a - \tilde{\Phi}^a \right|^2 + \frac{1}{t} |x^m - y^m|^2 + \\ + \frac{1}{s_1} |\partial_a W(y)|^2 + \frac{1}{\sqrt{s_1 s_2}} \partial_a \partial^b W(y) \Gamma^a \hat{\Gamma}_b + \frac{1}{s_2} |p_a|^2 + \frac{1}{s_2} |\nabla_m|^2 + \dots, \end{aligned} \quad (42)$$

where $\nabla_m = p_m + i A_m(x^l | y^a, p_a)$ and dots stand for gauge dependent terms which are irrelevant after taking the limit $s_1, s_2 \rightarrow 0$.

⁴ One could say that we have $N = 1$ SUSY Quantum mechanic in the directions orthogonal to D-brane and $N = \frac{1}{2}$ SUSY Quantum mechanic in the directions parallel to D-brane.

Let us define the symmetric trace "SymTr" following [33, 34] as the symmetrisation of the usual trace. Note that:

$$\text{Tr } e^{A+B+C} = \text{SymTr } e^{A+B+C} \quad (43)$$

Now substituting (42) into (22) and taking $\text{Tr}_{\mathcal{H}}$, one finds in the limit $t, s_1, s_2 \rightarrow 0$:

$$S_{RR} = Q_7 \int_{M^{(7)}} \text{SymTr} \left[\exp \left\{ -\frac{1}{2\pi} \left(\hat{F}_{(2)}(x^m) + [\hat{\nabla}_{(1)}(x^m), i_{\Phi}] + [i_{\Phi}, i_{\Phi}] \right) \right\} C_{RR}(x^m, \Phi^a) \right]_{top}, \quad (44)$$

where $M^{(7)}$ is the $D7$ -brane world-volume, $\hat{F}_{(2)} = [\hat{\nabla}_m, \hat{\nabla}_n] dx^m \wedge dx^n$, $\hat{\nabla}_{(1)} = \hat{\nabla}_m dx^m$ and i_{Φ} is defined by [34]:

$$i_{\Phi} C_{...a} dx^a = C_{...a} \hat{\Phi}^a.$$

The last two formulas contain $k \times k$ matrixes $\hat{A}$ and $\hat{\Phi}$ instead of $A(x|y, p)$ and $\tilde{\Phi}(x|y, p)$ as in the section 4. Note that the expression (44) coincides with S_{RR} considered in [34]. Thus, we observe that the massless excitations A and Φ on D -branes appear as Goldstone bosons due to the breaking of the symmetry generated by the algebra of symplectomorphisms down to $U(k)$ algebra.

8. It is easy to consider within this framework most general configuration of parallel $Dp - \overline{Dp}$ -brane systems with various p along the lines of [31]. As well it is not hard to consider deformations of the corresponding superconnections which describe non-parallel D -branes [31].

It would be interesting to apply our procedure to topological string theories, where the approximation of string non-commutativity structures by the algebra of differential operators is exact at least for some correlation functions. As well it would be interesting to generalize all our considerations to the case of the algebra of differential operators on the space of paths of space-time.

I am grateful to A.Gerasimov for very valuable discussions and S.Shatashvili for collaboration. As well I would like to thank organizers of the Tohwa International Symposium and especially Tada Tsukasa and Hajime Aoki for giving me an opportunity to present my results and for hospitality. This work was partially supported by RFBR 01-01-00548 and was commissioned by the AIP.

References

- [1] A.Polyakov, "Gauge Fields and Strings", (Heidelberg etc.: Springer-Verlag, 1987).
- [2] E. Witten, "Noncommutative Geometry And String Field Theory," Nucl. Phys. **B268** (1986) 235.
- [3] L.Rastelly, A.Sen and B.Zwiebach, "Half Strings, Projectors And Multiple D-branes In Vacuum String Field Theory", hep-th/0105058; "Boundary CFT Construction Of D-brane In Vacuum String Field Theory", hep-th/0105168
- [4] D.Gross and W.Taylor, "Split String Field Theory", hep-th/0105059.
- [5] E. Witten, "On Background Independent Open String Field Theory," hep-th/9208027, Phys. Rev. **D46** (1992) 5467; E. Witten, "Some Computations In Background Independent Open-String Field Theory," hep-th/9210065, Phys. Rev. **D47** (1993) 3405.

- [6] S. Shatashvili, "Comment On The Background Independent Open String Theory," hep-th/9303143, Phys. Lett. **B311** (1993) 83; S. Shatashvili, "On The Problems With Background Independence In String Theory," Preprint IASSNS-HEP-93/66, hep-th/9311177, Algebra and Anal. **v.6** (1994) 215.
- [7] A. Sen, "Stable Non-BPS States In String Theory," hep-th/9803194, JHEP **9806:007** (1998); A. Sen, "Stable Non-BPS Bound States Of BPS D-branes," hep-th/9805019, JHEP **9808:010** (1998); A. Sen, "Tachyon Condensation On The Brane Antibrane System," hep-th/9805170, JHEP **9808:012** (1998).
- [8] A. Gerasimov and S. Shatashvili, "On The Exact Tachyon Potential In Open String Field Theory," hep-th/0009103, JHEP **0010:034** (2000).
- [9] D. Kutasov, M. Marino and G. Moore, "Some Exact Results On Tachyon Condensation In String Field Theory", hep-th/0009148, JHEP **0010:045** (2000).
- [10] D. Kutasov, M. Marino and G. Moore, "Remarks On Tachyon Condensation In Superstring Field Theory", hep-th/0010108.
- [11] A. Gerasimov and S. Shatashvili, "Stringy Higgs Mechanism And The Fate Of Open Strings," hep-th/0011009, JHEP **0101:019** (2001).
- [12] M. Douglas, "Branes Within Branes," hep-th/9512077.
- [13] M. Green, J. Harvey and G. Moore, "I-Brane Inflow And Anomalous Couplings On D-Branes," hep-th/9605033, Calss. Quant. Grav., **14** (1997) 47.
- [14] M. Billio, B. Craps and F. Roose, "Ramond-Ramond Couplings Of Non-BPS D-branes," hep-th/9905157, JHEP **9906:033** (1999).
- [15] D. Freed and E. Witten, "Anomalies In String Theory With D-branes," hep-th/9907189.
- [16] J. Morales, C. Scrucca and M. Serone, "Anomalous Couplings For D-branes And O-planes," hep-th/9812071, Nucl. Phys. **B552** (1999) 291.
- [17] B.Stefanski, Nucl.Phys., **B548**, (1999) 275.
- [18] C. Kennedy and A. Wilkins, "Ramond-Ramond Couplings On Brane-Antibrane Systems," hep-th/9905195, Phys. Lett. **B464** (1999) 206.
- [19] K. Hori, "D-Branes, T Duality, And Index Theory," hep-th/9902102, Adv. Theor. Math. Phys. **3** (1999) 281.
- [20] P. Kraus and F. Larsen, "Boundary String Field Theory Of The $D\overline{D}$ System," hep-th/0012198, Phys. Rev. **D63** (2001) 106004.
- [21] T. Takayanagi, S. Terashima and T. Uesugi, "Brane-Antibrane Action From Boundary String Field Theory," hep-th/0012210, JHEP **0103:019** (2001).
- [22] K.Olesen and R.Szabo, Adv. Theor. Math. Phys., **3** (1999) 889.
- [23] Mohsen Alishahiha, Harald Ita and Yaron Oz, "*On Superconnections and the Tachyon Effective Action*," hep-th/0012222, Phys. Lett. **B503** (2001) 181-188.

- [24] S. F. Hassan and R. Minasian, "D-brane Couplings, RR Fields And Clifford Multiplication," hep-th/0008149.
- [25] R. Minasian and A. Tomasiello, "Variations On Stability," hep-th/0104041.
- [26] J. Schwarz and E. Witten, "Anomaly Analysis Of Brane-Antibrane Systems," hep-th/0103099, JHEP **0103:032** (2001).
- [27] E. Witten, "Overview Of K-Theory Applied To Strings," hep-th/0007175, Int. J. Mod. Phys. **A16** (2001) 693.
- [28] G. Moore and R. Minasian, "K-theory And Ramond-Ramond Charges," hep-th/9710230.
- [29] E. Witten, "D-branes And K-theory," hep-th/9810188, JHEP **9812:019** (1998).
- [30] D. Quillen, Topology **24** (1985) 89; V.Mathai and D.Quillen, Topology **25** (1986) 85.
- [31] E.Akhmedov, A.Gerasimov and S.Shatashvili, "On Unification of RR couplings", hep-th/0105228.
- [32] A. Gerasimov and S. Shatashvili, "Non-abelian Structures In Open String Theory," hep-th/0105245.
- [33] A. Tseytlin, "On Non-abelian Generalisation Of Born-Infeld Action In String Theory," hep-th/9701125, Nucl. Phys. **B501** (1997) 41.
- [34] R. Myers, "Dielectric-Branes," hep-th/9910053, JHEP **9912:022** (1999).
- [35] S. Donaldson, P. Kronheimer, "The Geometry Of Four-manifolds," Clarendon Press, Oxford, 1990, 440pp.
- [36] E. Witten, "Supersymmetry And Morse theory," J. Diff. Geom. **17** (1982) 661.
- [37] A. Alekseev, L. Faddeev and S. Shatashvili, "Quantization Of Symplectic Orbits Of Compact Lie Groups By Means Of The Functional Integral," J. Geom. Phys. 5:391-406 (1989);
A. Alekseev and S. Shatashvili, "From Geometric Quantization To Conformal Field Theory," Commun. Math. Phys. **128** (1990) 197;
A.Yu. Morozov, A. J. Niemi and K. Palo, Phys.Lett., **B271** (1991) 365.